

Preparing for your Colonoscopy Procedure

Thank you for choosing the Gastroenterology Associates at Mass General Hospital for your colonoscopy. We want to make sure your procedure goes as smoothly as possible. **Please read all the instructions in this packet at least 1 week before your colonoscopy.** All instructions must be followed or your colonoscopy may be cancelled. If you have any questions about your medicines or your prep instructions, call your MGH Gastroenterologist. If you have any other questions, call us at 617-726-7663.

Date and arrival time:					
Please Note: There is more than one location for MGH colonoscopy procedures. Your location is listed here.					
Procedure location:					
Scheduled procedure:					
Patient name:					
MGH Gastroenterologist name and phone number:					
call us at 617-726-7663.					

Please note: Your arrival time is different than the start time for your colonoscopy so you have time to get ready for your procedure.

- Most patients are ready for discharge within 3 hours of the scheduled arrival time but occasionally unforeseen events occur that result in delays. You will be informed of any potential delays when you arrive.
- Your escort should be available to meet you within 30 minutes after we call.



Plan ahead

Update your MGH registration information by calling 866-211-6588.
☐ If you have questions about the coverage for your procedures, please call and verify with your insurance company directly.
☐ Sign up for a Partners HealthCare Patient Gateway account if you do not have one. It will help with communicating with us. You will be able to see your test results in Patient Gateway within 2 weeks of the procedure.
Arrange for an adult escort to take you home after your procedure.
Read your prep instructions carefully. Plan to buy any items you will need to start your prep.
☐ If your procedure is scheduled at Charles River Plaza (165 Cambridge St) and you use CPAP, home oxygen or have an implantable cardiac defibrillator, call so your exam can be rescheduled in the Blake Building.
Please note that MGH policy requires that women, ages 11-55 years old have a pregnancy test prior to having any endoscopic procedure. When you arrive for your procedure, a registered nurse will screen you for the test and if needed, request that you provide a urine sample.
Read the information about the day of your procedure in this packet. It will tell you what to bring.

It is very important that you keep this appointment. **If you must cancel, please call us at least 5 business days before your appointment** by calling 617-726-7663. Calling ahead allows us to reschedule your appointment and give that slot to another patient. **If you cancel late, we may not be able to reschedule your appointment.**

For driving directions, please visit the MGH Parking and Visitor Information website at www.massgeneral.org/visit. If you are using GPS, please make sure to use the correct zip code.

For more information and frequently asked questions, please visit our website www.massgeneral.org/endoscopy.



MiraLAX Shopping List and Diet

This sheet tells you what you will need to do to clean your bowels so that your doctor can see polyps or other problems inside your colon. If your bowels are not clean, you may have to reschedule your test. If you have questions about this colonoscopy prep, call your MGH gastroenterologist.

Items v	vou	will	need	to	buy
	, 0 0-	,,			~ U - 1

Laxative pills. Either generic bisacodyl or Dulcolax®. You will need 2 pills.
Laxative powder. Either generic polyethylene glycol or MiraLAX®. You will need 238 grams or an 8.3 ounce bottle. Do not mix it with water until the afternoon before your procedure.
Gatorade, Crystal Light or water to mix with your laxative powder. Remember no red or orange liquids.
Simethicone anti-gas pills. Any brand is fine. Examples include Gas-X Mylanta Gas, or Maalox Anti-Gas.
Clear liquids that are not colored red, purple or orange. Clear liquid is any liquid that you can see through. Examples include water, tea, black coffee, clear broth, apple juice, white grape juice, sodas, sports drinks like Gatorade, and Jell-O.
Products to help with anal irritation including baby wipes and Vaseline. If
you have hemorrhoids you can use cream or Tucks pads.
■ Milk of magnesia. If you move your bowels 2 times a week or less or if you use a laxative more than 2 times a month, buy milk of magnesia.

Medications

- If you have diabetes, ask your primary care practitioner about changes in your diabetes medicine dose. If you take insulin, we usually recommend that you take ½ your normal dose on the day of the procedure.
- If you take blood thinners, we recommend you take them unless your MGH gastroenterology doctor told you to stop taking them.



MiraLAX Bowel Preparation Instructions

Five (5) days before your procedure:

If you take vitamins, iron pills, or liquid antacids, stop taking them 5 days before your procedure.

Two (2) days before your procedure:

- Don't eat popcorn, seeds, nuts, salad, corn, beans, peas, whole grain or whole wheat breads, raw fruits or raw vegetables: well-cooked fruits and vegetable are acceptable.
- -If you move your bowels 2 times a week or less or use a laxative more than 2 times a month, at **bedtime**, take 4 tablespoons of milk of magnesia.

One (1) day before your procedure:

Follow a clear liquid diet only. Clear liquid is any liquid that you can see through. Do not drink any red, purple or orange liquids. Examples of clear liquids include: Water, tea, black coffee, clear broth, apple juice, white grape juice, sodas, sports drinks like Gatorade, and Jell-O.

Remember! The day before your procedure, don't drink or eat:

- solid foods
- dairy products
- alcoholic drinks

4 pm (the day before your procedure)

• Take 2 bisacodyl pills with water.

6 pm (the day before your procedure)

- Mix the entire 238 gram container of MiraLAX powder with 64 ounces of liquid. You may use water, Gatorade or Crystal Light.
- Drink half of the laxative container.
- Start by drinking 8 ounces which is 1 cup of the laxative.
- Repeat this drink every 10 minutes until you have had about ½ of the total container of laxative.
- Keep drinking the laxative on schedule even though you may not start moving your bowels for 2 to 3 hours.

MiraLAX Bowel Preparation Instructions

(continued)

9 pm (the day before your procedure)

• Take 2 gas pills with 8 ounces of clear liquid.

10 pm (the day before your procedure)

• Take 2 gas pills with 8 ounces of clear liquid.

Day of your procedure:

If your arrival time is early in the morning, you will need to finish drinking the laxative in the early morning hours even if this means you will not get much sleep. Take your morning pills with clear liquids. If you stopped taking vitamins, iron pills, or liquid antacids, do not take them today.

6 hours before your scheduled arrival time:

- Drink the last half of the laxative.
- Drink 8 ounces which is 1 cup of the laxative.
- Repeat this drink every 10 minutes until the laxative is gone.
- You must drink ALL the laxative 4 hours before your arrival time.

2 hours before your scheduled arrival time:

- Stop drinking all liquids.
- Do not chew gum or eat any food.
- If you drink liquids, chew gum, eat hard candy or eat food within 2 hours of your procedure, it will be postponed or cancelled.

Comfort tips

- If you feel sick to your stomach, wait 30 minutes then start drinking smaller amounts.
- Try drinking the laxative through a straw.
- If you don't like the taste of the laxative, try chewing gum or sucking on hard candy in between drinks.
- Use baby wipes, Vaseline, or hemorrhoid cream if you get sore from moving your bowels.

If you have questions about these colonoscopy preparation instructions, call your MGH gastroenterologist.



The Day of Your Colonoscopy Procedure

Bring these things with you to your colonoscopy procedure
☐ Your photo identification.
☐ The name and phone number of your escort.
☐ You may wear your wedding rings but no other jewelry.

The day of your procedure

- The time for your appointment is earlier than the time your procedure will start so you can get ready.
- Before the procedure, we will review the procedure with you and ask you to sign a consent form. (see last page)
- Most procedures take about 3 hours. We make every effort to keep on time, but sometimes there are delays.
- We will call your escort 30 minutes before you are ready to leave.

After the procedure

- Most people need to rest at home for the remainder of the day. Don't drive or operate any machines on the day of your procedure. Avoid making any important decisions. Avoid drinking alcohol.
- You can go back to eating as you normally do right away.
- You will get a letter in the mail with your test results within 2 weeks after your colonoscopy. If you have a Partners HealthCare Patient Gateway account, you can also see your results there.

Remember

- You cannot drive after your procedure.
- We will have to cancel your procedure if you do not have an adult escort to meet you in the endoscopy unit and bring you home.
- Your escort should be able to pick you up 30 minutes after we call them.

For any questions about this information call 617-726-7663.





PATIENT CONSENT TO PROCEDURE

	PATIENT:					
	UNIT NO:					
	PROCEDURE: Colonoscopy with possible biopsy or polypectomy					
	☐ Right ☐ Left ☐ Both Sides ☑ Not applicable					
	My doctor has told me and I understand what procedure/surgery I am having risks (like drug reactions, bleeding, infection, and complications from receiving no guarantee of results. My doctor has also explained what might happen to can make instead of having this done, (including choosing no treatment) and something else. I understand that with any procedure, problems could come to me how he/she prevents infections related to my health. The following additional contents of the contents of t	g blood or blood components), and that there is me if I don't have this procedure, other choices I what can happen to me if I choose to do up that we did not expect. My provider explained				
\supset	Colonoscopy is a procedure for examination of the colon through which biopsies may be taken and polyps removed. Although colonoscopy is generally safe, some infrequent and possibly serious complications may occur. The most common problem is a reaction to one of the drugs given for the purpose of sedation. Some of the complications of conscious sedation include low blood pressure, low oxygen levels, and changes in heart rhythm. More serious complications of colonoscopy are hemorrhage and perforation, which may require hospitalization, blood transfusion, or surgery. These complications are more frequent following polyp removal, but serious problems usually occur in less than one percent of patients. An additional risk of colonoscopy is the possibility that a polyp, lesion or even a cancer could be missed during the examination.					
\supset						
	If procedural sedation will be used during this procedure to control my pain, I risks. These risks include difficulty breathing that may require breathing supple common side effects are nausea and vomiting. In rare cases, there can be all heart). Lastly, I may have pain, even after using these medications.	ort and decreased blood pressure. The most				
\supset	My doctor and/or his/her associate will perform my procedure/surgery. I understand that Massachusetts General means that resident doctors, doctors in a medical fellowship (fellows) and study care professions receive training here, and may take part in my procedure/surgether to perform my procedure/surgery. My doctor or an attending designed procedure/surgery, although other medical professionals may perform some attending designee deems appropriate.	Hospital (MGH) is a teaching hospital. This dents in medical, nursing and related health gery. A team of medical professionals will work ee will be present for all the critical parts of the				
	I understand that this procedure/surgery may have educational or scientific value. The hospital may photograph, videotape or record my procedure/surgery for educational, research, quality and other healthcare operations purposes. Any informati used for these purposes will not identify me.					
	I understand that blood or other samples removed during this procedure may also may be used by MGH, its partners, or affiliates for research, education an					
	I have had the chance to ask questions about the risks, benefits and alternative answers I received. I consent to this procedure/surgery.	es to this procedure/surgery. I am happy with the				
	Date Time AM/PM Signature (patient/health care agent/guardian/family member) (If patient's con	sent cannot be obtained, indicate reason above.)				
J	I attest that I discussed all relevant aspects of this procedure/surgery, including with alternative approaches, with the patient, and answered his/her questions.	-				
	Date Time AM/PM					
	Signature (Physician/Licensed	Practitioner)				